

Five Decades of the Gravitational Weyl anomaly

M. J. Duff

Blackett Laboratory, Imperial College London & Mathematical Institute, Oxford

Gravity: past, present and future
UBC & PITP
September 2017

Classical Weyl invariance

- Classically, Weyl invariance

$$S(g, \phi) = S(g', \phi')$$

under

$$g'_{\mu\nu}(x) = \Omega(x)^2 g_{\mu\nu}(x) \quad \Phi'(x) = \Omega(x)^\alpha \Phi(x)$$

implies

$$g^{\mu\nu} T_{\mu\nu} = 0$$

- To accommodate fermions $e'^a_\mu(x) = \Omega(x)e_\mu^a(x)$ where $g_{\mu\nu}(x) = e_\mu^a(x)e_{\nu a}(x)$

Examples

- Conformally coupled scalar, $\phi'(x) = \Omega\phi(x)$, Penrose

$$S[\phi] = \int d^4x \sqrt{-g} \left(-\frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - \frac{1}{6} R \phi^2 \right)$$

- Massless fermion, $\psi'(x) = \Omega^{-3/2}(x)\psi(x)$, Dirac

$$S[\psi] = \int d^4x e (\bar{\psi} \gamma^\mu \nabla_\mu \psi)$$

- Electromagnetic field, $A'_\mu(x) = A_\mu(x)$, Maxwell

$$S[A] = \int d^4x \sqrt{-g} \left(-\frac{1}{4} g^{\mu\rho} g^{\nu\sigma} F_{\mu\nu} F_{\rho\sigma} \right)$$

Quantum Weyl anomalies

- But in the quantum theory

$$g^{\mu\nu} \langle T_{\mu\nu} \rangle \neq 0$$

Over the period 1973-2012 this Weyl anomaly has found a variety of applications in quantum gravity, black hole physics, inflationary cosmology, string theory and statistical mechanics.

- Note that generic curved space

$$g^{\mu\nu} T_{\mu\nu}$$

not associated with a Noether current

Recall flat space ancestry 1970

- For example $SO(D, 2)$ in the case of flat Minkowski space.
Coleman and Jackiw Callan, Coleman and Jackiw
- More generally, for D-dimensional spacetimes admitting conformal Killing vectors $\xi_{\mu}^i(x)$

$$\nabla_{\mu}\xi_{\nu}^i + \nabla_{\nu}\xi_{\mu}^i = \frac{2}{D}g_{\mu\nu}\nabla^{\rho}\xi_{\rho}^i$$

there is a classically conserved dilatation current

$$J^{i\nu} = \xi_{\mu}^i T^{\mu\nu}$$

- Anomaly appears in the quantum theory

$$\nabla_{\nu} \langle J^{i\nu} \rangle = \frac{1}{D} \nabla^{\rho} \xi_{\rho}^i g^{\mu\nu} \langle T_{\mu\nu} \rangle \neq 0$$

but this is not an anomaly in local Weyl symmetry

$$g'_{\mu\nu}(x) = \Omega(x)^2 g_{\mu\nu}(x)$$

Timeline 1973

- Corrections to graviton propagator from closed loops of spin $s = 0, 1/2, 1$ using dimensional regularization.
Capper Duff and Halpern $s = 1$
Capper and Duff $s = 1/2$
Geist et al $s = 0$
- Discovery of the Weyl anomaly using dimensional regularization
Capper and Duff

Timeline 1974

- I first announced the existence of gravitational Weyl anomalies at The First Oxford Quantum Gravity Conference, organised by **Isham, Penrose, Sciama**, and held at the Rutherford Laboratory in February 1974
- Unfortunately, the announcement was somewhat overshadowed because **Hawking** chose the same conference to reveal to an unsuspecting world his result that black holes evaporate!
- Ironically, **Christensen, Fulling** were subsequently to link the Hawking effect and the trace anomaly.

Timeline 1976

- Non-local effective lagrangian for trace anomaly **Deser, Duff and Isham** By general covariance and dimensional analysis, it must take the following form:

- For $D=2$,

$$g^{\alpha\beta} \langle T_{\alpha\beta} \rangle = aR$$

- For $D=4$,

$$g^{\alpha\beta} \langle T_{\alpha\beta} \rangle = \alpha R^2 + \beta R_{\mu\nu} R^{\mu\nu} + \gamma R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} + \delta \square R + c F_{\mu\nu}^a F^{\mu\nu a}$$

where $a, \alpha, \beta, \gamma, \delta$ and c are constants.

- For $D = 6$,

$$g^{\alpha\beta} \langle T_{\alpha\beta} \rangle = (\text{curvature})^3$$

and so on.

- At one-loop, and ignoring boundary terms, there is no anomaly for D odd.

King's College London 1976



The heat kernel

- Zeta functions, heat kernels and anomalies

Christensen

Dowker

Hawking

Barvinsky et al

The heat kernel

- Classical action

$$S_0 = \int d^d x \frac{1}{2}(\Phi, \Delta\Phi)$$

where Δ is a conformally invariant d-dimensional operator.

- The one-loop effective action is given by

$$S_1 = \ln[\det\Delta]^{-1/2}$$

- Its kernel $F(x, y, \rho)$ obeys the heat equation

$$\frac{\partial}{\partial\rho} F(x, y, \rho) + \Delta F(x, y, \rho) = 0$$

with the initial conditions

$$F(x, y, 0) = \delta(x, y)$$

The heat kernel

- One can express F as

$$\begin{aligned} F(x, y, \rho) &= \sum_n e^{-\rho\Delta} \phi_n(x) \phi_n(y) \\ &= \sum_n e^{-\rho\lambda_n} \phi_n(x) \phi_n(y) \end{aligned}$$

where ϕ_n are the eigenfunctions of Δ with eigenvalues λ_n :

$$\Delta\phi_n = \lambda_n\phi_n$$

normalized according to

$$\int d^d x \sqrt{g(x)} \phi_n(x) \phi_m(x) = \delta_{mn}$$

b_4 coefficients

- The one-loop action may thus be written as

$$S_1 = \int d\rho d^d x \rho^{-1} \sqrt{g}(x) A(x, \rho)$$

where $A(x, \rho) = F(x, x, \rho)$. $A(x, \rho)$ obeys an asymptotic expansion, valid for small ρ ,

$$A(x, \rho) \sim \sum_n B_n(x) \rho^{n - \frac{d}{2}}$$

where

$$B_n = \int d^d x \sqrt{g} b_n(x) \quad (1)$$

Zeta functions

- The **Schwinger-DeWitt** coefficients b_n are scalar polynomials, which are of order n in derivatives of the metric. In $d = 4$, for example, when Δ is the conformally invariant Laplacian acting on scalars:

$$\Delta = -\square + \frac{1}{6}R \quad \text{Penrose}$$

$$g^{\mu\nu} T_{\mu\nu} = b_4 = \frac{1}{2880\pi^2} [R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} - R_{\mu\nu} R^{\mu\nu} + 30\square R]$$

- Furthermore,

$$B_4 = n_0 + \zeta(0)$$

where n_0 is the number of zero modes and

$$\zeta(s) = \sum_n \lambda_n^{-s}$$

is defined only over the non-zero eigenvalues of Δ .

Timeline 1976

- Asymptotic safety
Weinberg
- Point splitting regularization
Christensen Duncan
- More anomaly coefficients
Dowker and Critchley Duncan
- Vacuum energy in two dimensions
Davies and Fulling
- Particle creation
Wald
- Robertson-Walker and applications to cosmology
Birrell, Bunch, Christensen, Davies, Fulling
Hartle et al
- Black holes
Davies, Fulling, Unruh

Timeline 1977

- CFTs and the a and c coefficients
Duff
- Trace anomalies and the Hawking effect
Christensen and Fulling

Conformal Field Theories (CFT)

- Weyl anomalies appear in their most pristine form when CFTs are coupled to an external gravitational field. In this case

$$\mathcal{A} = g^{\mu\nu} \langle T_{\mu\nu} \rangle = \frac{1}{(4\pi)^2} (cF - aG)$$

where F is the square of the Weyl tensor:

$$F = C_{\mu\nu\rho\sigma} C^{\mu\nu\rho\sigma} = R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} - 2R_{\mu\nu} R^{\mu\nu} + \frac{1}{3}R^2,$$

G is proportional to the Euler density:

$$G = R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} - 4R_{\mu\nu} R^{\mu\nu} + R^2,$$

- Note no R^2 term.
- We ignore $\square R$ terms whose coefficient can be adjusted to any value by adding the finite counterterm

$$\int d^4x \sqrt{g} R^2.$$

Central charges c and a

- In the CFT a and c are the central charges given in terms of the field content by

$$\bar{a} \equiv 720a = 2N_0 + 11N_{1/2} + 124N_1$$

$$\bar{c} \equiv 720c = 6N_0 + 18N_{1/2} + 72N_1$$

where N_s are the number of fields of spin s .

- In the notation of [Duff 1977](#)

$$(4\pi)^2 b = c \quad (4\pi)^2 b' = -a$$

Central charges c and a

- The story of the Weyl anomaly for CFTs is thus the story of the central charges c and a . The ratio is given by

$$\frac{a}{c} = \frac{(2N_0 + 11N_{1/2} + 124N_1)}{(6N_0 + 18N_{1/2} + 72N_1)}$$

and by inspection we can read off see the inequalities

$$\frac{31}{18} \geq \frac{a}{c} \geq \frac{1}{3}$$

where the upper and lower bounds are saturated by a single vector and a single scalar respectively.

Euler number

- When $F - G$ vanishes, anomaly reduces to

$$\mathcal{A} = A \frac{1}{32\pi^2} R^{*\mu\nu\rho\sigma} R^*_{\mu\nu\rho\sigma}$$

where

$$360A = \bar{c} - \bar{a} = 4N_0 + 7N_{1/2} - 52N_1$$

so that in Euclidean signature

$$\int d^4x \sqrt{g} g^{\mu\nu} T_{\mu\nu} = A\chi(M^4)$$

where $\chi(M^4)$ is the Euler number of spacetime.

Timeline 1978

- Conformal (and axial) anomalies for arbitrary spin
Christensen,Duff
- Conformal anomalies for interacting theories: QED, ϕ^4 etc
Drummond
Shore
Hathrell

Arbitrary spin

- Calculate b_4 for arbitrary (n, m) reps of Lorentz group, then physical anomaly given by

$$A = A(n, m) + A(n - 1, m - 1) - 2A(n - 1/2, m - 1/2)$$

so in total

$$A_{total} = 4N_0 + 7N_{1/2} - 52N_1 - 233N_{3/2} + 848N_2$$

where N_s are the number of fields of spin s .

- The b_4 coefficient for chiral reps $(1/2, 0)$ $(1, 0)$ etc also involve R^*R unless we add $(0, 1/2)$ $(0, 1)$ etc

Timeline 1980

- Anomaly-driven inflation
Starobinsky
Grishchuk, Zeldovich
Vilenkin
- p -forms and inequivalent anomalies
Duff, van Nieuwenhuizen
Grisaru et al
Siegel
- The path-integral approach to anomalies
Fujikawa
Bastianelli, van Nieuwenhuizen
Nicolai, Townsend

WARNING

- THE FOLLOWING CONTENT CONTAINS REFERENCES TO SUPERSYMMETRY WHICH MAY OFFEND SOME MEMBERS OF THE AUDIENCE
- VIEWER DISCRETION IS ADVISED

Central charges c and a

- In the supersymmetric case we have the values and bounds given below. Remarkably, these bounds continue to hold true when the CFT is interacting **Maldacena**

Fields	a	c	Bounds
$\mathcal{N} = 0$ spin 0	$1/360$	$1/120$	$31/18 \geq a/c \geq 1/3$
$\mathcal{N} = 0$ spin 1/2	$11/720$	$1/40$	
$\mathcal{N} = 0$ spin 1	$31/180$	$1/10$	
$\mathcal{N} = 1$ chiral multiplet	$1/48$	$1/24$	$3/2 \geq a/c \geq 1/2$
$\mathcal{N} = 1$ vector multiplet	$3/16$	$1/8$	
$\mathcal{N} = 2$ hyper multiplet	$1/24$	$1/12$	$5/4 \geq a/c \geq 1/2$
$\mathcal{N} = 2$ vector multiplet	$5/24$	$1/6$	
$\mathcal{N} = 4$ vector multiplet	$1/4$	$1/4$	$a/c = 1$

Table: The central charges a and c for supersymmetric CFTs

Timeline 1977

- Trace of stress tensor $T^\mu{}_\mu$
Divergence of axial current $\partial_\mu J^5_\mu$
Gamma trace of spinor current $\gamma^\mu S_\mu$
form a supermultiplet
- and so, therefore, do the anomalies!
Ferrara,Zumino

Timeline 1981

- When we allow for a cosmological constant Λ the anomaly is

$$A\chi + BV$$

where V is the volume. We find

$$B = 6N_0 + 18N_{1/2} + 72N_1 - 822N_{3/2} + 3132N_2 \quad (2)$$

Moreover in gauged supergravity

$$e^2 = G\Lambda$$

and B also determines the Yang-Mills beta-function.

- This yields vanishing β -function in gauged $N > 4$ supergravity [Christensen,Duff,Gibbons,Rocek](#)
- Spin sum rules

$$\sum_{\lambda} (-1)^{2\lambda} \lambda^k = 0$$

for $N > k$ [Curtwright Christensen,Duff](#)

- Critical dimensions for bosonic and super strings
Polyakov

Bosonic string

- In the first quantized theory of the bosonic string, one starts with a Euclidean functional integral

$$e^{-\Gamma} = \int \frac{D\gamma DX}{\text{Vol}(\text{Diff})} e^{-S[\gamma, X]}$$

where

$$S[\gamma, X] = \frac{1}{4\pi\alpha'} \int d^2\xi \sqrt{\gamma} \gamma^{ij} \partial_i X^\mu \partial_j X^\nu \eta_{\mu\nu}$$

- As shown by **Polyakov**, the Weyl anomaly in the worldsheet stress tensor is given by

$$\gamma^{ij} \langle T_{ij} \rangle = \frac{1}{24\pi} (D - 26) R(\gamma)$$

D is the contribution of the scalars while the -26 arises from the diffeomorphism ghosts that must be introduced into the functional integral.

Fermionic string

- In the case of the fermionic string, the result is

$$\gamma^{ij} \langle T_{ij} \rangle = \frac{1}{16\pi} (D - 10) R(\gamma)$$

- Thus the critical dimensions $D = 26$ and $D = 10$ correspond to the preservation of the two dimensional Weyl invariance $\gamma_{ij} \rightarrow \Omega^2(\xi)\gamma_{ij}$.

Spacetime Einstein equations from worldsheet anomaly



$$S[\gamma, X] = \frac{1}{4\pi\alpha'} \int d^2\xi \sqrt{\gamma} \gamma^{ij} \partial_i X^\mu \partial_j X^\nu g_{\mu\nu}$$
$$\beta(g)_{\mu\nu} = R_{\mu\nu} + \dots$$

vanishing anomaly implies Einstein equations! Callan,
Friedan, Perry

Timeline 1983

- Conformal anomaly and W-Z consistency (no R^2)
[Bonora et al](#)
- Anomaly in conformal supergravity $N = 1, 2, 3, 4$

$$S = \int d^4x \sqrt{-g} C^{\mu\nu\rho\sigma} C_{\mu\nu\rho\sigma} + \dots$$

Vanishes for $N = 4$ [Fradkin and Tseytlin](#)

Timeline 1984

- Local version of effective action
Riegert

Local action

- Conformal operators

$$\sqrt{g}\Delta_d = \sqrt{g'}\Delta'_d$$

$$\Delta_2 = \square$$

$$\Delta_4 \equiv \square^2 + 2R^{\mu\nu}\nabla_\mu\nabla_\nu + \frac{1}{3}(\nabla^\mu R)\nabla_\mu - \frac{2}{3}R\square$$

Riegert

- Subsequent work by
Osborn et al Antoniadis, Mazur and Mottola Barvinsky et al
- Local action

$$S_{anom} = \frac{b}{2} \int d^4x \sqrt{g} F \phi - \frac{b'}{2} \int d^4x \sqrt{g} [\phi \Delta_4 \phi - (G - \frac{2}{3} \square R) \phi]$$

Timeline 1985

- Conformal invariants
Fefferman, Graham

- The c -theorem
Zamolodchikov

Timeline 1988

- c -theorem and/or a -theorem in four dimensions?

Cardy

Osborn

Capelli et al

Shore

Shapere

Antoniadis et al

- Geometric classification of conformal anomalies in arbitrary dimensions
Deser, Schwimmer

Timeline 1998

- The holographic Weyl anomaly
Henningson,Skenderis
Imbimbo
Graham
Bastianelli
Manvelyan
Fukuma
- Einstein manifolds and the a and c coefficients
Gubser,Martelli

Holography

- A distinguished coordinate system, boundary at $\rho = 0$

$$G_{MN}dx^M dx^N = \frac{L_{d+1}^2}{4} \rho^{-2} d\rho d\rho + \rho^{-1} g_{\mu\nu} dx^\mu dx^\nu$$

- The effective action may be written

$$S_B = \int d\rho d^d x \rho^{-1} \sqrt{g}(x) B(x, \rho)$$

where the specific form of $B(x, \rho)$ depends on initial action.

$$B(x, \rho) \sim \sum_n b_n(x) \rho^{n - \frac{d}{2}}$$

- Formal similarity with **Schwinger-DeWitt** coefficients, indeed $\mathcal{A} \sim b_4$ same for N=4 Yang-Mills but not in general.

Timeline 2000

- Anomaly-driven inflation revived
Hawking et al
Hamada
Nojiri
Shapiro
de Paula Netto, Pelinson, Shapiro, Starobinsky
- a and c and corrections to Newton's law
Duff and Liu
- Anomalies and entropy bounds
Nojiri et al

Corrections to Newton's law

- In my 1972 PhD thesis, at the suggestion of Abdus Salam, I calculated one-loop CFT corrections to Newton's law (Schwarzschild solution)

$$V(r) = \frac{G_4 M}{r} \left(1 + \frac{8cG_4}{3\pi r^2} \right),$$

where G_4 is the four-dimensional Newton's constant and c is a purely numerical coefficient. In fact it turned out to be the c coefficient in the Weyl anomaly

N=4 Yang-Mills

- A particularly important example of a CFT is provided by $\mathcal{N} = 4$ super Yang-Mills with gauge group $U(N)$, for which

$$(N_1, N_{1/2}, N_0) = (N^2, 4N^2, 6N^2)$$

Then

$$a = c = \frac{N^2}{4}$$

and hence

$$\mathcal{A} = \frac{c}{(4\pi)^2} \left(2R_{\mu\nu}R^{\mu\nu} - \frac{2}{3}R^2 \right) = \frac{N^2}{32\pi^2} \left(R_{\mu\nu}R^{\mu\nu} - \frac{1}{3}R^2 \right)$$

- The contribution of a single $\mathcal{N} = 4$ $U(N)$ Yang-Mills CFT is

$$V(r) = \frac{G_4 M}{r} \left(1 + \frac{2N^2 G_4}{3\pi r^2} \right).$$

Randall-Sundrum

- Now fast-forward to 1999 when **Randall and Sundrum** proposed that our four-dimensional world is a 3-brane embedded in an infinite five-dimensional universe. They showed that there is an r^{-3} correction coming from the massive Kaluza-Klein modes

$$V(r) = \frac{G_4 M}{r} \left(1 + \frac{2L_5^2}{3r^2} \right).$$

where L_5 is the radius of AdS_5 .

- Superficially, our 4D quantum correction seems unrelated to their 5D classical one.
- But through the miracle of AdS/CFT

$$N^2 = \frac{\pi L_5^3}{2G_5} \quad G_4 = \frac{2G_5}{L_5}$$

the two are in fact equivalent. **Duff and Liu**

Timeline 2001

- a and c and the graviton mass
Dilkes et al
Aharony
- Weyl cohomology revisited
Mazur and Mottola

- Anomalies as an infra-red diagnostic; IR free or interacting?
Intriligator

- Macroscopic effects of the quantum trace anomaly
Mottola et al
Gianotti et al

- Anomalies and the hierarchy problem
Meissner

Timeline 2008

- Viscosity bounds
Buchel et al
- Conformal collider physics
Hofman and Maldacena
- Weyl invariance and mass
Waldron et al

- Entanglement Entropy
Nishioka
- Log corrections to black hole entropy
Cai
Solodukin
Sen et al

- Holographic c-theorems in arbitrary dimensions
Myers et al
- Generalized mirror symmetry and trace anomalies
Duff et al
- Vanish without a trace
Duff et al

M-theory on X^7

- We consider compactification of $(\mathcal{N} = 1, D = 11)$ supergravity on a 7-manifold X^7 with betti numbers $(b_0, b_1, b_2, b_3, b_3, b_2, b_1, b_0)$ and define a generalized mirror symmetry

$$(b_0, b_1, b_2, b_3) \rightarrow (b_0, b_1, b_2 - \rho/2, b_3 + \rho/2)$$

under which

$$\rho(X^7) \equiv 7b_0 - 5b_1 + 3b_2 - b_3$$

changes sign

$$\rho \rightarrow -\rho$$

- The massless sectors of these compactifications have

$$f = 4(b_0 + b_1 + b_2 + b_3)$$

degrees of freedom.

- Generalized self-mirror theories are defined to be those for which $\rho = 0$

M-theory on X^7

- In backgrounds for which $F - G$ vanishes, the Weyl anomaly reduces to

$$T = A \frac{1}{32\pi^2} R^{*\mu\nu\rho\sigma} R^*_{\mu\nu\rho\sigma} \quad (3)$$

where

$$A = 2(c - a) \quad (4)$$

so that in Euclidean signature

$$\int d^4x \sqrt{g} T = A \chi(M^4) \quad (5)$$

where $\chi(M^4)$ is the Euler number of spacetime.

Anomalies

	<i>Field</i>	<i>f</i>	ΔA	$360A$	$360A'$	X^7
g_{MN}	$g_{\mu\nu}$	2	-3	848	-232	b_0
	\mathcal{A}_μ	2	0	-52	-52	b_1
	\mathcal{A}	1	0	4	4	$-b_1 + b_3$
ψ_M	ψ_μ	2	1	-233	127	$b_0 + b_1$
	χ	2	0	7	7	$b_2 + b_3$
A_{MNP}	$A_{\mu\nu\rho}$	0	2	-720	0	b_0
	$A_{\mu\nu}$	1	-1	364	4	b_1
	A_μ	2	0	-52	-52	b_2
	A	1	0	4	4	b_3

total ΔA

0

total A

$-\rho/24$

total A'

$-\rho/24$

Vanish without a trace!

- Remarkably, we find that the anomalous trace depends on ρ

$$A = -\frac{1}{24}\rho(X^7)$$

So the anomaly flips sign under generalized mirror symmetry and vanishes for generalized self-mirror theories. For $X^{(8-\mathcal{N})} \times T^{(\mathcal{N}-1)}$ with $\mathcal{N} \geq 3$ the anomaly vanishes identically.

Duff and Ferrara

- Equally remarkable is that we get the same answer for the total trace using the numbers of [Grisaru et al.](#)

Four curious supergravities

- Of particular interest are the four cases

$$(b_0, b_1, b_2, b_3) = (1, \mathcal{N} - 1, 3\mathcal{N} - 3, 4\mathcal{N} + 3)$$

with $\mathcal{N} = 1, 2, 4, 8$, namely the four “curious” supergravities, discussed in [Duff and Ferrara](#) which enjoy some remarkable properties.

$\mathcal{N} = 1$, 7 WZ multiplets, $f = 32$,

$\mathcal{N} = 2$, 3 vector multiplets, 4 hypermultiplets, $f = 64$,

$\mathcal{N} = 4$, 6 vector multiplets, $f = 128$,

$\mathcal{N} = 8$, $f = 256$.

O, H, C, R theories

<i>Field</i>	360A	O	H	C	R
$g_{\mu\nu}$	848	1	1	1	1
B_μ	-52	7	6	0	0
S	4	28	16	10	7
ψ_μ	-233	8	4	2	1
χ	7	56	28	14	7
$A_{\mu\nu\rho}$	-720	1	1	1	1
$A_{\mu\nu}$	364	7	3	1	0
A_μ	-52	21	6	4	0
A	4	35	19	11	7
		$A = 0$	$A = 0$	$A = 0$	$A = 0$

Table: Vanishing anomaly in **O, H, C, R** theories.

Fano plane




Figure: The Fano plane has seven points and seven lines (the circle counts as a line) with three points on every line and three lines through every point. The truncation from 7 lines to 3 to 1 to 0 corresponds to the truncation from $N=8$ to $N=4$ to $N=2$ to $N=1$.

Type IIA

- In the case of $(\mathcal{N} = 1, D = 11)$ on $X^6 \times S^1$, or equivalently (Type IIA, $D=10$) on X^6 ,

$$A = -\frac{1}{24}\chi(X^6)$$

and so in Euclidean signature

$$\int d^4x \sqrt{g} g_{\mu\nu} \langle T^{\mu\nu} \rangle = -\frac{1}{24}\chi(M^4)\chi(X^6) = -\frac{1}{24}\chi(M^{10})$$

where $\chi(M^4)$ is the Euler number of spacetime.

Timeline 2011

- Models for particle physics
't Hooft
- Renormalization group and Weyl anomalies
Percacci
- A four-dimensional a-theorem
Komargodski et al
Luty et al
Elvang et al

- Gravitational anomalies and thermal Hall effect in topological insulators
Stone
- A one-loop test of quantum gravity
Bhattacharyya et al

Timeline 2015

- Holographic c-theorems in arbitrary dimensions
Stone
- A one-loop test of quantum supergravity
Bhattacharyya et al
- Anomalies and conformal manifolds
Gomis
- More on boundary terms in the anomaly
Fursaev
Solodukhin

- The semi-classical stress-energy tensor in a Schwarzschild background
Bardeen

Acknowledgements

- Grateful to Leron Borston, Steve Christensen, Stanley Deser, Marc Grisaru, Emil Mottola, Ashoke Sen for discussions.
- Thanks to the organizers for the invitation.